

References: Semicircular Canal Dehiscence: Lessons for Collaboration, in partnership with the American Academy of Audiology

Presenter: Rebekah F. Cunningham, PhD; Devin L. McCaslin, PhD

- Meiklejohn DA, Corrales CE, Boldt BM, Sharon JD, Yeom KW, Carey JP, Blevins NH. Pediatric Semicircular Canal Dehiscence: Radiographic and Histologic Prevalence, With Clinical Correlation. *Otol Neurotol*. 2015 Sep;36(8):1383-9. doi: 10.1097/MAO.0000000000000811. PMID: 26164444.
- Merchant SN, Rosowski JJ. Conductive hearing loss caused by third-window lesions of the inner ear. *Otol Neurotol*. 2008 Apr;29(3):282-9. doi: 10.1097/mao.0b013e318161ab24. PMID: 18223508; PMCID: PMC2577191.
- Chen EY, Paladin A, Phillips G, Raske M, Vega L, Peterson D, Sie KC. Semicircular canal dehiscence in the pediatric population. *Int J Pediatr Otorhinolaryngol*. 2009 Feb;73(2):321-7. doi: 10.1016/j.ijporl.2008.10.027. Epub 2008 Dec 16. PMID: 19091428.
- Minor LB, Solomon D, Zinreich JS, Zee DS. Sound- and/or pressure-induced vertigo due to bone dehiscence of the superior semicircular canal. *Arch Otolaryngol Head Neck Surg*. 1998 Mar;124(3):249-58. doi: 10.1001/archotol.124.3.249. PMID: 9525507.
- Carey JP, Minor LB, Nager GT. Dehiscence or thinning of bone overlying the superior semicircular canal in a temporal bone survey. *Arch Otolaryngol Head Neck Surg*. 2000 Feb;126(2):137-47. doi: 10.1001/archotol.126.2.137. PMID: 10680863.