

Audiological and Vestibular Care of Patients with Parkinson's Disease (PD)

Melissa Newell, AuD

Reference List

Acarer, A., Karapolat, H., Celebisoy, N., Ozgen, G., Colakoglu, Z (2015). Is customized vestibular rehabilitation effective in patients with Parkinson's? *NeuroRehabilitation*, 37(2), 255-262.
doi:10.3233/NRE-151258

Allum, J. H., Tang, K. S., Carpenter, M. G., Oude Nijhuis, L. B., & Bloem, B. R. (2011). Review of first trial responses in balance control: influence of vestibular loss and Parkinson's disease. *Human movement science*, 30(2), 279–295. <https://doi.org/10.1016/j.humov.2010.11.009>

Bertolini, G., Wicki, A., Baumann, C. R., Straumann, D., & Palla, A. (2015). Impaired tilt perception in Parkinson's disease: a central vestibular integration failure. *PloS One*, 10(4), e0124253.
<https://doi.org/10.1371/journal.pone.0124253>

Beylergil, S. B., Ozinga, S., Walker, M. F., McIntyre, C. C., & Shaikh, A. G. (2019). Vestibular heading perception in Parkinson's disease. *Progress in Brain Research*, 249, 307–319.
<https://doi.org/10.1016/bs.pbr.2019.03.034>

Bohnen, N. I., Roytman, S., Griggs, A., David, S. M., Beaulieu, M. L., & Müller, M. L. T. M. (2022). Decreased vestibular efficacy contributes to abnormal balance in Parkinson's disease. *Journal of the Neurological Sciences*, 440, 120357. <https://doi.org/10.1016/j.jns.2022.120357>

Cui, W., Duan, Z., & Feng, J. (2022). Assessment of vestibular-evoked myogenic potentials in Parkinson's Disease: a systematic review and meta-analysis. *Brain Sciences*, 12(7), 956.
<https://doi.org/10.3390/brainsci12070956>

Giladi, N., Mirelman, A., Thaler, A., Bar-Shira, A., Gurevich, T., & Orr-Urtreger, A. (2011). Fighting the risk of developing Parkinson's disease; clinical counseling for first degree relatives of patients with Parkinson's disease. *Journal of the Neurological Sciences*, 310(1-2), 17-20.
<https://doi.org/10.1016/j.jns.2011.06.005>

Gökay, N. Y., Gündüz, B., Söke, F., & Karamert, R. (2020). Evaluation of efferent auditory system and hearing quality in Parkinson's Disease: is the difficulty in speech understanding in complex listening conditions related to neural degeneration or aging? *Journal of Speech, Language, and Hearing Research*, 64(1), 263-271. https://doi.org/10.1044/2020_JSLHR-20-00337

Hawkins, K. E., Holden, J., Chiarovano, E., Lewis, S. J. G., Curthoys, I. S., & MacDougall, H. G. (2025). Vestibular evoked myogenic potentials (VEMPs) in Parkinson's Disease patients with monopolar deep-brain stimulation. *Signals*, 6(1), 10. <https://doi.org/10.3390/signals6010010>

Keyser, K. D., Letter, M. D., Groote, E. D., Santens, P., Talsma, D., Botteldooren, D., & Bockstael, A. (2019). Systematic audiological assessment of Auditory functioning in

patients with Parkinson's Disease. *Journal of Speech, Language, and Hearing Research*, 62(12), 4564-4577. doi: 10.1044/2019_JSLHR-H-19-0097.

Kia, M., Mirian, M. S., Soori, S., Saedi, S., Arasteh, E., Faramarzi, M. H., Chinchani, A., Lee, S., Luczak, A., & McKeown, M. J. (2025). Koopman-based linearization of preparatory EEG dynamics in Parkinson's disease during galvanic vestibular stimulation. *Frontiers in Human Neuroscience*, 19, 1566566. <https://doi.org/10.3389/fnhum.2025.1566566>

Leme, M. S., Sanches, S. G. G., & Carvalho, R. M. M. (2022). Peripheral hearing in Parkinson's disease: a systematic review. *International Journal of Audiology*, 62(9), 805–813. <https://doi.org/10.1080/14992027.2022.2109073>

Manso, A., Ganança, M. M., & Caovilla, H. H. (2016). Vestibular rehabilitation with visual stimuli in peripheral vestibular disorders. *Brazilian Journal of Otorhinolaryngology*, 82(2), 232. <https://doi.org/10.1016/j.bjorl.2015.05.019>

Menon, A., Vigneswaran, M., Zhang, T., Sreenivasan, V., Kim, C., & McKeown, M. J. (2025). Individualising galvanic vestibular stimulation further improves visuomotor performance in Parkinson's Disease. *Bioengineering*, 12(5), 523. <https://doi.org/10.3390/bioengineering12050523>

Morozetti, P. G., Ganança, C. F., & Chiari, B. M. (2011). Comparison of different protocols for vestibular rehabilitation in patients with peripheral vestibular disorders. *Jornal da Sociedade Brasileira de Fonoaudiologia*, 23(1), 44–50. <https://doi.org/10.1590/s2179-64912011000100011>

National Institute on Aging. *Parkinson's Disease: Causes, Symptoms, and Treatments*. <https://www.nia.nih.gov/health/parkinsons-disease/parkinsons-disease-causes-symptoms-and-treatments>

Ninja Nerd (viewed May 5, 2025); Youtube.com. *Parkinson's Disease/Clinical Presentation/ Part 1*. https://youtu.be/KWVJBg6SCoY?si=onJSC9PVHiy_rlsK

Ninja Nerd (viewed May 5, 2025); Youtube.com. *Parkinson's Disease/Causes and Pathophysiology/ Part 2*. <https://youtu.be/rFoc4ACFehQ?si=rv-8Qzr76WwjCRZ5>

Neilson, L.E., Reavis, K.M., Wiedrick, J., Scott, G.D. (2024). Hearing loss, incident Parkinson Disease, and treatment with hearing aids. *JAMA Neurology*, 81(12), 1295–1303. doi:10.1001/jamaneurol.2024.3568

Pastor, M. A., Day, B. L., & Marsden, C. D. (1993). Vestibular induced postural responses in Parkinson's disease. *Brain: A Journal of Neurology*, 116 (Pt 5), 1177–1190. <https://doi.org/10.1093/brain/116.5.1177>

Ray Chaudhuri, K., Poplawska-Domaszewicz, K., Limbachiya, N., Qamar, M., Batzu, L., Podlewska, A., & Ade, K. (2024). Vestibular neurostimulation for Parkinson's Disease: a novel device-aided non-invasive therapeutic option. *Journal of Personalized Medicine*, 14(9), 933. <https://doi.org/10.3390/jpm14090933>

Shalash, A. S., Hassan, D. M., Elrassas, H. H., Salama, M. M., Méndez-Hernández, E., Salas-Pacheco, J. M., & Arias-Carrión, O. (2017). Auditory- and vestibular-evoked potentials correlate with motor and non-motor features of Parkinson's Disease. *Frontiers in Neurology*, 8, 55. <https://doi.org/10.3389/fneur.2017.00055>

Smith, P. F. (2018). Vestibular functions and Parkinson's Disease. *Frontiers in Neurology*, 9, 1085. <https://doi.org/10.3389/fneur.2018.01085>

Vitale, C., Marcelli, V., Abate, T., Pianese, A., Allocca, R., Moccia, M., Spina, E., Barone, P., Santangelo, G., & Cavaliere, M. (2016). Speech discrimination is impaired in Parkinsonian patients: expanding the audiologic findings of Parkinson's disease. *Parkinsonism & Related Disorders*, 22, S138-S143. <https://doi.org/10.1016/j.parkreldis.2015.09.040>

Wilkinson, D., Podlewska, A., Banducci, S. E., Pellat-Higgins, T., Slade, M., Bodani, M., Sakel, M., Smith, L., LeWitt, P., & Ade, K. (2019). Caloric vestibular stimulation for the management of motor and non-motor symptoms in Parkinson's disease: intention-to-treat data. *Data in Brief*, 25, 104228. <https://doi.org/10.1016/j.dib.2019.104228>

Xie, W. Y., Duan, W. X., Chen, Y., Tao, M. X., Li, H. X., Gao, F., Yin, J. Y., Yan, J. H., Wang, F., Mao, C. J., Shen, Y., & Liu, C. F. (2025). The impact of bright light therapy on Parkinson's disease: a pilot study using vestibular-evoked myogenic potentials. *Parkinsonism & Related Disorders*, 134, 107776. <https://doi.org/10.1016/j.parkreldis.2025.107776>

Yakubovich, S., Israeli-Korn, S., Halperin, O., Yahalom, G., Hassin-Baer, S., & Zaidel, A. (2020). Visual self-motion cues are impaired yet overweighted during visual-vestibular integration in Parkinson's disease. *Brain Communications*, 2(1), fcaa035. <https://doi.org/10.1093/braincomms/fcaa035>

Zeigelboim, B. S., Klagenberg, K. F., Teive, H. A., Munhoz, R. P., & Martins-Bassetto, J. (2009). Vestibular rehabilitation: clinical benefits to patients with Parkinson's disease. *Arquivos de Neuro-Psiquiatria*, 67(2A), 219–223. <https://doi.org/10.1590/s0004-282x2009000200009>